



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS AND SOLUTIONS.

B. F. FINKEL, CHAIRMAN OF THE COMMITTEE.

PROBLEMS FOR SOLUTION.

ALGEBRA.

396. Proposed by H. E. TREFETHEN, Colby College.Show that $1 + \frac{1}{3} - \frac{1}{5} - \frac{1}{7} + \frac{1}{9} + \frac{1}{11} - \cdots = \sqrt{2}(1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots)$.

GEOMETRY.

424. Proposed by H. E. TREFETHEN, Colby College.In a given triangle ABC , determine by geometric demonstration the point O so that the sum of the distances, $AO + BO + CO$, shall be a minimum.**425. Proposed by V. M. SPUNAR, Chicago, Ill.**Find the ratio of the areas, A_1 and A_2 , of the parabolas formed by projectiles whose ranges are the same and whose angles of projection are complements of each other.

CALCULUS.

345. Proposed by C. N. SCHMALL, New York City.

"Of all the quadrilaterals which can be formed from four given sides, that which is inscriptible in a circle has the greatest area."

[From GOURSAT-HEDRICK, *Math. Anal.*, p. 133, ex. 5.]*Proposer's Remark.*—Prove this, and furthermore show that when only three sides are known, the length of the fourth side of the maximum quadrilateral is the root of a cubic equation.**346. Proposed by C. N. SCHMALL, New York City.**

Given the height of an inclined plane, to find its length so that a given force, acting on a given mass in a direction parallel to the plane, may draw it up in the shortest time.

Note.—This question has a practical application in the case of a truck, of known height, upon which a mass is to be loaded by means of skids.

MECHANICS.

280. Proposed by C. N. SCHMALL, New York City.Given the distance d between two smooth hooks in the same horizontal line. Show that the shortest string which can form a catenary, with these hooks for points of support, is de , where e is the base of the Napierian system of logarithms.**281. Proposed by C. N. SCHMALL, New York City.** ABC is a triangle inscribed in a circle, center O , and L, M, N , are the centers of gravity of the sectors AOB, BOC, COA . Show that

$$\frac{AB}{OL} + \frac{BC}{OM} + \frac{CA}{ON} = 3\pi.$$

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

197. Proposed by E. T. BELL, Seattle, Wash.Show that in the expansion of $\frac{1 + z + z^2 + \cdots + z^{p-1}}{(1 - z)^{p-1}} - 1$, where p is a prime, the coefficients of the various powers of z are divisible by p . [EISENSTEIN, *Crelle*, t. 27, p. 282.]